

# Transverse structure of the nucleon

## Part 2: Theory background

---

Alessandro Bacchetta



# DIS and structure functions


---

# Deep inelastic scattering

---


$$\ell(l) + N(P) \rightarrow \ell(l') + h(P_h) + X,$$

$$x_B = \frac{Q^2}{2 P \cdot q}, \quad y = \frac{P \cdot q}{P \cdot l}, \quad z_h = \frac{P \cdot P_h}{P \cdot q}.$$



# Inclusive DIS


---



$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2 M W^{\mu\nu}(q, P, S)$$

# Inclusive DIS

---




$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2 M W^{\mu\nu}(q, P, S)$$

*Single-photon-exchange approximation*

# Inclusive DIS


---



$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2 M W^{\mu\nu}(q, P, S)$$

*Single-photon-exchange approximation*

$$\begin{aligned} L_{\mu\nu} &= \sum_{\lambda'_e} \left( \bar{u}(l', \lambda'_e) \gamma_\mu u(l, \lambda_e) \right)^* \left( \bar{u}(l', \lambda'_e) \gamma_\nu u(l, \lambda_e) \right) \\ &= -Q^2 g_{\mu\nu} + 2 \left( l_\mu l'_\nu + l'_\mu l_\nu \right) + 2i \lambda_e \epsilon_{\mu\nu\rho\sigma} l^\rho l'^\sigma \\ &= \frac{2Q^2}{y^2} \left[ - \left( 1 - y + \frac{y^2}{2} \right) g_{\perp\mu\nu} + 2(1-y) \hat{t}_\mu \hat{t}_\nu \right. \\ &\quad + 2(1-y) \left( \hat{l}_\mu \hat{l}_\nu + \frac{1}{2} g_{\perp\mu\nu} \right) \\ &\quad \left. - i \lambda_e y \left( 1 - \frac{y}{2} \right) \epsilon_{\perp\mu\nu} - i \lambda_e y \sqrt{1-y} \hat{t}_{[\mu} \epsilon_{\perp\nu]\rho} \hat{l}^\rho + \dots \right] \end{aligned}$$



# Lepton tensor

---

$$l^\mu = \left( \frac{(2-y)Q}{2y}, \frac{\sqrt{1-y}Q}{y}, 0, \frac{Q}{2} \right),$$
$$l'^\mu = \left( \frac{(2-y)Q}{2y}, \frac{\sqrt{1-y}Q}{y}, 0, -\frac{Q}{2} \right)$$

# Lepton tensor

---

$$l^\mu = \left( \frac{(2-y)Q}{2y}, \frac{\sqrt{1-y}Q}{y}, 0, \frac{Q}{2} \right),$$
$$l'^\mu = \left( \frac{(2-y)Q}{2y}, \frac{\sqrt{1-y}Q}{y}, 0, -\frac{Q}{2} \right)$$

$$L_{\mu\nu}(\lambda_e = 0) = \frac{2Q^2}{y^2} \begin{pmatrix} 2(1-y) & \dots & 0 & 0 \\ \dots & \left(1-y + \frac{y^2}{2}\right) + (1-y) & 0 & 0 \\ 0 & 0 & \left(1-y + \frac{y^2}{2}\right) - (1-y) & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# Lepton tensor

---

$$l^\mu = \left( \frac{(2-y)Q}{2y}, \frac{\sqrt{1-y}Q}{y}, 0, \frac{Q}{2} \right),$$

$$l'^\mu = \left( \frac{(2-y)Q}{2y}, \frac{\sqrt{1-y}Q}{y}, 0, -\frac{Q}{2} \right)$$

$$L_{\mu\nu}(\lambda_e = 0) = \frac{2Q^2}{y^2} \begin{pmatrix} 2(1-y) & \dots & 0 & 0 \\ \dots & \left(1-y + \frac{y^2}{2}\right) + (1-y) & 0 & 0 \\ 0 & 0 & \left(1-y + \frac{y^2}{2}\right) - (1-y) & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\epsilon_x^\mu = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad \epsilon_y^\mu = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad \epsilon_L^\mu = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

# Helicity vs transversity / circular vs linear

---

$$\chi_+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$\chi_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$\chi_\uparrow = \frac{1}{\sqrt{2}}(\chi_+ + e^{i\phi_s}\chi_-) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ e^{i\phi_s} \end{pmatrix}$$

$$\chi_\downarrow = \frac{1}{\sqrt{2}}(\chi_+ - e^{i(\phi_s+\pi)}\chi_-) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ e^{i(\phi_s+\pi)} \end{pmatrix}$$

# Helicity vs transversity / circular vs linear

---

$$\chi_+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$\chi_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$\chi_\uparrow = \frac{1}{\sqrt{2}}(\chi_+ + e^{i\phi_s}\chi_-) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ e^{i\phi_s} \end{pmatrix}$$

$$\chi_\downarrow = \frac{1}{\sqrt{2}}(\chi_+ - e^{i(\phi_s+\pi)}\chi_-) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ e^{i(\phi_s+\pi)} \end{pmatrix}$$

$$\epsilon_+^\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ i \\ 0 \end{pmatrix},$$

$$\epsilon_-^\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -i \\ 0 \end{pmatrix},$$

$$\epsilon_x^\mu = \frac{1}{\sqrt{2}}(\epsilon_+^\mu + \epsilon_-^\mu) = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix},$$

$$\epsilon_y^\mu = \frac{i}{\sqrt{2}}(\epsilon_+^\mu - \epsilon_-^\mu) = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

# Helicity conservation

---



$$\bar{u}_+(-p_z)\gamma^\mu u_+(p_z) \sim (0, 1, i, 0)$$

$$\bar{u}^\uparrow(-p_z)\gamma^\mu u^\uparrow(p_z) \sim (0, \cos\phi_s, -\sin\phi_s, 0)$$

$$\bar{u}_-(-p_z)\gamma^\mu u_+(p_z) = 0$$

$$\bar{u}^\downarrow(-p_z)\gamma^\mu u^\uparrow(p_z) = 0$$

# Structure functions

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$

# Structure functions

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$


$$2MW^{\mu\nu} = \frac{1}{x} \left[ -g_{\perp}^{\mu\nu} F_{UU,T} + \hat{t}^\mu \hat{t}^\nu F_{UU,L} + iS_L \epsilon_{\perp}^{\mu\nu} F_{LL} - i\hat{t}^{[\mu} \epsilon_{\perp}^{\nu]\rho} S_\rho F_{LT}^{\cos \phi_S} \right]$$

# Structure functions

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$

$$2MW^{\mu\nu} = \frac{1}{x} \left[ -g_{\perp}^{\mu\nu} F_{UU,T} + \hat{t}^{\mu} \hat{t}^{\nu} F_{UU,L} + iS_L \epsilon_{\perp}^{\mu\nu} F_{LL} - i\hat{t}^{[\mu} \epsilon_{\perp}^{\nu]\rho} S_{\rho} F_{LT}^{\cos \phi_S} \right]$$




# Structure functions

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$

$$2MW^{\mu\nu} = \frac{1}{x} \left[ -g_{\perp}^{\mu\nu} F_{UU,T} + \hat{t}^\mu \hat{t}^\nu F_{UU,L} + iS_L \epsilon_{\perp}^{\mu\nu} F_{LL} - i\hat{t}^{[\mu} \epsilon_{\perp}^{\nu]\rho} S_\rho F_{LT}^{\cos \phi_S} \right]$$



$$\begin{aligned} \frac{d\sigma}{dx_B dy d\phi_S} = & \frac{2\alpha^2}{x_B y Q^2} \left\{ \left(1 - y + \frac{y^2}{2}\right) F_{UU,T} + (1 - y) F_{UU,L} + S_L \lambda_e y \left(1 - \frac{y}{2}\right) F_{LL} \right. \\ & \left. + |S_T| \lambda_e y \sqrt{1 - y} \cos \phi_S F_{LT}^{\cos \phi_s} \right\} \end{aligned}$$

# Semi-inclusive DIS

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$

# Semi-inclusive DIS

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$

$$\frac{2E_h d^6\sigma}{d^3P_h dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S, P_h)$$

# Semi-inclusive DIS

---

$$\frac{d^3\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S)$$

$$\frac{2E_h d^6\sigma}{d^3P_h dx_B dy d\phi_S} = \frac{\alpha^2 y}{2 Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S, P_h)$$

$$\frac{d^6\sigma}{dx_B dy dz_h d\phi_S d\phi_h dP_{h\perp}^2} = \frac{\alpha^2 y}{2 z_h Q^4} L_{\mu\nu}(l, l', \lambda_e) 2MW^{\mu\nu}(q, P, S, P_h)$$

# Structure functions

---

$$\begin{aligned}
& \frac{d\sigma}{dx dy d\phi_S dz d\phi_h dP_{h\perp}^2} \\
&= \frac{\alpha^2}{x y Q^2} \frac{y^2}{2(1-\varepsilon)} \left\{ \right. \\
&\quad F_{UU,T} + \varepsilon F_{UU,L} + \sqrt{2\varepsilon(1+\varepsilon)} \cos \phi_h F_{UU}^{\cos \phi_h} + \varepsilon \cos(2\phi_h) F_{UU}^{\cos 2\phi_h} \\
&\quad + \lambda_e \sqrt{2\varepsilon(1-\varepsilon)} \sin \phi_h F_{LU}^{\sin \phi_h} + S_L \left[ \sqrt{2\varepsilon(1+\varepsilon)} \sin \phi_h F_{UL}^{\sin \phi_h} + \varepsilon \sin(2\phi_h) F_{UL}^{\sin 2\phi_h} \right] \\
&\quad + S_L \lambda_e \left[ \sqrt{1-\varepsilon^2} F_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} \cos \phi_h F_{LL}^{\cos \phi_h} \right] \\
&\quad + S_T \left[ \sin(\phi_h - \phi_S) \left( F_{UT,T}^{\sin(\phi_h - \phi_S)} + \varepsilon F_{UT,L}^{\sin(\phi_h - \phi_S)} \right) + \varepsilon \sin(\phi_h + \phi_S) F_{UT}^{\sin(\phi_h + \phi_S)} \right. \\
&\quad + \varepsilon \sin(3\phi_h - \phi_S) F_{UT}^{\sin(3\phi_h - \phi_S)} + \sqrt{2\varepsilon(1+\varepsilon)} \sin \phi_S F_{UT}^{\sin \phi_S} \\
&\quad \left. + \sqrt{2\varepsilon(1+\varepsilon)} \sin(2\phi_h - \phi_S) F_{UT}^{\sin(2\phi_h - \phi_S)} \right] + S_T \lambda_e \left[ \sqrt{1-\varepsilon^2} \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)} \right. \\
&\quad \left. + \sqrt{2\varepsilon(1-\varepsilon)} \cos \phi_S F_{LT}^{\cos \phi_S} + \sqrt{2\varepsilon(1-\varepsilon)} \cos(2\phi_h - \phi_S) F_{LT}^{\cos(2\phi_h - \phi_S)} \right] \left. \right\}
\end{aligned}$$

see e.g. AB, Diehl, Goeke, Metz, Mulders, Schlegel, JHEP093 (07)

# Structure functions

---

$$\begin{aligned}
& \frac{d\sigma}{dx dy d\phi_S dz d\phi_h dP_{h\perp}^2} F_{UU,T}(x, z, P_{h\perp}^2, Q^2) \\
&= \frac{\alpha^2}{x y Q^2} \frac{y^2}{2(1-\varepsilon)} \left\{ F_{UU,T} + \varepsilon F_{UU,L} + \sqrt{2\varepsilon(1+\varepsilon)} \cos \phi_h F_{UU}^{\cos \phi_h} + \varepsilon \cos(2\phi_h) F_{UU}^{\cos 2\phi_h} \right. \\
&\quad + \lambda_e \sqrt{2\varepsilon(1-\varepsilon)} \sin \phi_h F_{LU}^{\sin \phi_h} + S_L \left[ \sqrt{2\varepsilon(1+\varepsilon)} \sin \phi_h F_{UL}^{\sin \phi_h} + \varepsilon \sin(2\phi_h) F_{UL}^{\sin 2\phi_h} \right] \\
&\quad + S_L \lambda_e \left[ \sqrt{1-\varepsilon^2} F_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} \cos \phi_h F_{LL}^{\cos \phi_h} \right] \\
&\quad + S_T \left[ \sin(\phi_h - \phi_S) \left( F_{UT,T}^{\sin(\phi_h - \phi_S)} + \varepsilon F_{UT,L}^{\sin(\phi_h - \phi_S)} \right) + \varepsilon \sin(\phi_h + \phi_S) F_{UT}^{\sin(\phi_h + \phi_S)} \right. \\
&\quad + \varepsilon \sin(3\phi_h - \phi_S) F_{UT}^{\sin(3\phi_h - \phi_S)} + \sqrt{2\varepsilon(1+\varepsilon)} \sin \phi_S F_{UT}^{\sin \phi_S} \\
&\quad \left. + \sqrt{2\varepsilon(1+\varepsilon)} \sin(2\phi_h - \phi_S) F_{UT}^{\sin(2\phi_h - \phi_S)} \right] + S_T \lambda_e \left[ \sqrt{1-\varepsilon^2} \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)} \right. \\
&\quad \left. + \sqrt{2\varepsilon(1-\varepsilon)} \cos \phi_S F_{LT}^{\cos \phi_S} + \sqrt{2\varepsilon(1-\varepsilon)} \cos(2\phi_h - \phi_S) F_{LT}^{\cos(2\phi_h - \phi_S)} \right] \}
\end{aligned}$$

see e.g. AB, Diehl, Goeke, Metz, Mulders, Schlegel, JHEP093 (07)

# Light-cone coordinates

---

Light-cone vectors will be indicated as

$$a^\mu = [a^-, a^+, \mathbf{a}_T] = \left[ \frac{a^0 - a^3}{\sqrt{2}}, \frac{a^0 + a^3}{\sqrt{2}}, a^1, a^2 \right]. \quad (1)$$

The dot-product in light-cone components is

$$\mathbf{a} \cdot \mathbf{b} = a^+ b^- + a^- b^+ - \mathbf{a}_T \cdot \mathbf{b}_T \quad (2)$$

The light-cone decomposition of a vector can be written in a Lorentz covariant fashion using two light-like vectors  $n_+$  and  $n_-$  satisfying  $n_\pm^2 = 0$  and  $n_+ \cdot n_- = 1$  and promoting  $\mathbf{a}_T$  to a four-vector  $a_T^\mu = [0, 0, \mathbf{a}_T]$  so that

$$a^\mu = a^+ n_+^\mu + a^- n_-^\mu + a_T^\mu, \quad (3)$$

where

$$a^+ = a \cdot n_-, \quad a^- = a \cdot n_+, \quad a_T \cdot n_+ = a_T \cdot n_- = 0. \quad (4)$$



Note that

$$a_T \cdot b_T = -\mathbf{a}_T \cdot \mathbf{b}_T \quad (5)$$

# Light-cone coordinates


---

$Q^2$  much higher than any other scalar product



# Correlation functions in DIS


---



$$2MW^{\mu\nu}(q, P, S) \approx \sum_q e_q^2 \frac{1}{2} \text{Tr} [\Phi(x_B, S) \gamma^\mu \gamma^+ \gamma^\nu].$$

# Correlation functions in DIS

---




$$2MW^{\mu\nu}(q, P, S) \approx \sum_q e_q^2 \frac{1}{2} \text{Tr} [\Phi(x_B, S) \gamma^\mu \gamma^+ \gamma^\nu].$$

$$\begin{aligned} \Phi_{ij}(x, S) &= \int d^2\mathbf{p}_T dp^- \left. \Phi_{ij}(p, P, S) \right|_{p^+ = xP^+} \\ &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) U_{[0, \xi]} \psi_i(\xi) | P, S \rangle \Big|_{\xi^+ = \xi_T = 0} \end{aligned}$$

# Correlation functions in DIS

---



$$2M W^{\mu\nu}(q, P, S) \approx \sum_q e_q^2 \frac{1}{2} \text{Tr} [\Phi(x_B, S) \gamma^\mu \gamma^+ \gamma^\nu].$$


$$\begin{aligned} \Phi_{ij}(x, S) &= \int d^2 \mathbf{p}_T dp^- \left. \Phi_{ij}(p, P, S) \right|_{p^+ = x P^+} \\ &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) U_{[0, \xi]} \psi_i(\xi) | P, S \rangle \Big|_{\xi^+ = \xi_T = 0} \end{aligned}$$

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4 \xi e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) U_{[0, \xi]} \psi_i(\xi) | P, S \rangle$$

# Correlation functions in SIDIS


---

$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{I} \left[ \text{Tr}(\Phi(x_B, \mathbf{p}_T, S) \gamma^\mu \Delta(z_h, \mathbf{k}_T) \gamma^\nu) \right]$$



# Correlation functions in SIDIS


$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{I} \left[ \text{Tr}(\Phi(x_B, \mathbf{p}_T, S) \gamma^\mu \Delta(z_h, \mathbf{k}_T) \gamma^\nu) \right]$$



$$\mathcal{I}[\dots] \equiv \int d^2\mathbf{p}_T d^2\mathbf{k}_T \delta^{(2)}(\mathbf{p}_T + \mathbf{q}_T - \mathbf{k}_T) [\dots] = \int d^2\mathbf{p}_T d^2\mathbf{k}_T \delta^{(2)}\left(\mathbf{p}_T - \frac{\mathbf{P}_{h\perp}}{z} - \mathbf{k}_T\right) [\dots]$$

# Correlation functions in SIDIS

$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{I} \left[ \text{Tr}(\Phi(x_B, \mathbf{p}_T, S) \gamma^\mu \Delta(z_h, \mathbf{k}_T) \gamma^\nu) \right]$$



$$\mathcal{I}[\dots] \equiv \int d^2\mathbf{p}_T d^2\mathbf{k}_T \delta^{(2)}(\mathbf{p}_T + \mathbf{q}_T - \mathbf{k}_T) [\dots] = \int d^2\mathbf{p}_T d^2\mathbf{k}_T \delta^{(2)}\left(\mathbf{p}_T - \frac{\mathbf{P}_{h\perp}}{z} - \mathbf{k}_T\right) [\dots]$$

*Only at low transverse momentum*

$$P_{h\perp}^2 \ll Q^2$$

# Integrated vs unintegrated correlators

---

$$\begin{aligned}\Phi_{ij}(x, S) &= \int d^2\mathbf{p}_T \Phi_{ij}(x, \mathbf{p}_T) \\ &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) U_{[0,\xi]} \psi_i(\xi) | P, S \rangle \Big|_{\xi^+ = \xi_T = 0}\end{aligned}$$

# Integrated vs unintegrated correlators

---

$$\begin{aligned}\Phi_{ij}(x, S) &= \int d^2\mathbf{p}_T \Phi_{ij}(x, \mathbf{p}_T) \\ &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) U_{[0, \xi]} \psi_i(\xi) | P, S \rangle \Big|_{\xi^+ = \boldsymbol{\xi}_T = 0}\end{aligned}$$

$$\begin{aligned}\Phi_{ij}(x, \mathbf{p}_T, S) &= \int dp^- \Phi(p, P, S) \Big|_{p^+ = xP^+} \\ &= \int \frac{d\xi^- d^2\boldsymbol{\xi}_T}{(2\pi)^3} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) U_{[0, \xi]} \psi_i(\xi) | P, S \rangle \Big|_{\xi^+ = 0}\end{aligned}$$

# Correlation functions

---

$p, P, S$

# Correlation functions

---

$p, P, S$

$1, \gamma_5, \gamma^\mu, \gamma^\mu \gamma_5, i\sigma^{\mu\nu} \gamma_5$

# Correlation functions

---

$$p,\,P,\,S$$

$$\mathbf{1},\,\gamma_5,\,\gamma^\mu,\,\gamma^\mu\gamma_5,\,i\sigma^{\mu\nu}\gamma_5\qquad\qquad\qquad \sigma^{\mu\nu}\equiv\frac{i}{2}\left[\gamma^\mu,\gamma^\nu\right].$$

# Correlation functions

---

$$p, P, S$$

$$1, \gamma_5, \gamma^\mu, \gamma^\mu \gamma_5, i\sigma^{\mu\nu} \gamma_5 \quad \sigma^{\mu\nu} \equiv \frac{i}{2} [\gamma^\mu, \gamma^\nu].$$

Hermiticity:  $\Phi(p, P, S) = \gamma^0 \Phi^\dagger(p, P, S) \gamma^0,$  (1a)

parity:  $\Phi(p, P, S) = \gamma^0 \Phi(\tilde{p}, \tilde{P}, -\tilde{S}) \gamma^0$  (1b)

# Correlation functions

---

$$p, P, S$$

$$1, \gamma_5, \gamma^\mu, \gamma^\mu \gamma_5, i\sigma^{\mu\nu} \gamma_5 \quad \sigma^{\mu\nu} \equiv \frac{i}{2} [\gamma^\mu, \gamma^\nu].$$

Hermiticity:  $\Phi(p, P, S) = \gamma^0 \Phi^\dagger(p, P, S) \gamma^0,$  (1a)

parity:  $\Phi(p, P, S) = \gamma^0 \Phi(\tilde{p}, \tilde{P}, -\tilde{S}) \gamma^0$  (1b)

$$\tilde{p}^\nu = \delta^{\nu\mu} p_\mu$$

# Correlation functions

---

$$p, P, S$$

$$\mathbf{1}, \gamma_5, \gamma^\mu, \gamma^\mu \gamma_5, i\sigma^{\mu\nu} \gamma_5 \quad \sigma^{\mu\nu} \equiv \frac{i}{2} [\gamma^\mu, \gamma^\nu].$$

Hermiticity:  $\Phi(p, P, S) = \gamma^0 \Phi^\dagger(p, P, S) \gamma^0,$  (1a)

parity:  $\Phi(p, P, S) = \gamma^0 \Phi(\tilde{p}, \tilde{P}, -\tilde{S}) \gamma^0$  (1b)

$$\tilde{p}^\nu = \delta^{\nu\mu} p_\mu$$

For an unpolarized target, the most general decomposition is

$$\Phi(p, P) = M A_1 \mathbf{1} + A_2 \not{P} + A_3 \not{p} + \frac{A_4}{M} \sigma_{\mu\nu} P^\mu p^\nu$$

where the amplitudes  $A_i$  are real scalar functions  $A_i = A_i(p \cdot P, p^2)$  with dimension  $1/[m]^4.$

# Parton distribution functions

---

# Parton distribution functions

---

If we keep only the leading terms in  $1/P^+$  (leading twist)

$$\Phi(p, P) \approx P^+ (A_2 + x A_3) \not{\epsilon}_+ + P^+ \frac{i}{2M} [\not{\epsilon}_+, \not{p}_T] A_4,$$

# Parton distribution functions

---

If we keep only the leading terms in  $1/P^+$  (leading twist)

$$\Phi(p, P) \approx P^+ (A_2 + x A_3) \not{\epsilon}_+ + P^+ \frac{i}{2M} [\not{\epsilon}_+, \not{p}_T] A_4,$$

$$\Phi(x, p_T) \equiv \int dp^- \Phi(p, P) = \frac{1}{2} \left\{ f_1 \not{\epsilon}_+ + i h_1^\perp \frac{[\not{p}_T, \not{\epsilon}_+]}{2M} \right\}.$$

# Parton distribution functions

---

If we keep only the leading terms in  $1/P^+$  (leading twist)

$$\Phi(p, P) \approx P^+ (A_2 + x A_3) \not{\epsilon}_+ + P^+ \frac{i}{2M} [\not{\epsilon}_+, \not{p}_T] A_4,$$

$$\Phi(x, p_T) \equiv \int dp^- \Phi(p, P) = \frac{1}{2} \left\{ f_1 \not{\epsilon}_+ + i h_1^\perp \frac{[\not{p}_T, \not{\epsilon}_+]}{2M} \right\}.$$

Here we introduced the parton distribution functions

$$f_1(x, p_T^2) = 2P^+ \int dp^- (A_2 + x A_3), \quad h_1^\perp(x, p_T^2) = 2P^+ \int dp^- (-A_4).$$

# Parton distribution functions

---

$$\Phi^{[\Gamma]} = \frac{1}{2} \text{Tr}[\Phi \Gamma]$$

$$\Phi^{[\gamma^+]} = f_1(x, p_T^2) ,$$

$$\Phi^{[i\sigma^\alpha + \gamma_5]} = -\frac{\epsilon_T^{\alpha\rho} p_{T\rho}}{M} h_1^\perp(x, p_T^2)$$

# Collinear PDFs and TMDs

---

$$\begin{aligned}\Phi(x, p_T) = & \frac{1}{2} \left\{ f_1 \not{\epsilon}_+ - f_{1T}^\perp \frac{\epsilon_T^{\rho\sigma} p_{T\rho} S_{T\sigma}}{M} \not{\epsilon}_+ + S_L g_{1L} \gamma_5 \not{\epsilon}_+ - g_{1T} \frac{p_T \cdot S_T}{M} \gamma_5 \not{\epsilon}_+ \right. \\ & \left. + h_{1T} \frac{[\not{S}_T, \not{\epsilon}_+] \gamma_5}{2} + h_{1s}^\perp \frac{[\not{p}_T, \not{\epsilon}_+] \gamma_5}{2M} + i h_1^\perp \frac{[\not{p}_T, \not{\epsilon}_+]}{2M} \right\}\end{aligned}$$

# Collinear PDFs and TMDs

---

$$\Phi(x) = \frac{1}{2} \left\{ f_1 \not{\epsilon}_+ + S_L g_1 \gamma_5 \not{\epsilon}_+ + h_1 \frac{[\not{S}_T, \not{\epsilon}_+] \gamma_5}{2} \right\}$$

$$\begin{aligned} \Phi(x, p_T) = & \frac{1}{2} \left\{ f_1 \not{\epsilon}_+ - f_{1T}^\perp \frac{\epsilon_T^{\rho\sigma} p_{T\rho} S_{T\sigma}}{M} \not{\epsilon}_+ + S_L g_{1L} \gamma_5 \not{\epsilon}_+ - g_{1T} \frac{p_T \cdot S_T}{M} \gamma_5 \not{\epsilon}_+ \right. \\ & \left. + h_{1T} \frac{[\not{S}_T, \not{\epsilon}_+] \gamma_5}{2} + h_{1s}^\perp \frac{[\not{p}_T, \not{\epsilon}_+] \gamma_5}{2M} + i h_1^\perp \frac{[\not{p}_T, \not{\epsilon}_+]}{2M} \right\} \end{aligned}$$

# Dirac matrices: an unusual representation

---

$$\gamma^0 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad \gamma^3 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix},$$

$$\gamma^1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad \gamma^2 = \begin{pmatrix} 0 & i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & -i & 0 \end{pmatrix},$$

$$\gamma_5 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

# Good/Bad and Right/Left projectors

---

$$\mathcal{P}^+ = \gamma^- \gamma^+ / 2,$$

$$\mathcal{P}^- = \gamma^+ \gamma^- / 2,$$

$$\mathcal{P}_R = (1 + \gamma_5) / 2,$$

$$\mathcal{P}_L = (1 - \gamma_5) / 2$$

# Good/Bad and Right/Left projectors

---

$$\mathcal{P}^+ = \gamma^- \gamma^+/2,$$

$$\mathcal{P}^- = \gamma^+ \gamma^-/2,$$

$$\mathcal{P}_R = (1 + \gamma_5)/2,$$

$$\mathcal{P}_L = (1 - \gamma_5)/2$$

$$\mathcal{P}_R \mathcal{P}^+ = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\mathcal{P}_L \mathcal{P}^+ = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\mathcal{P}_R \mathcal{P}^- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$\mathcal{P}_L \mathcal{P}^- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# Good/Bad and Right/Left projectors

---

$$\mathcal{P}^+ = \gamma^- \gamma^+/2,$$

$$\mathcal{P}^- = \gamma^+ \gamma^-/2,$$

$$\mathcal{P}_R = (1 + \gamma_5)/2,$$

$$\mathcal{P}_L = (1 - \gamma_5)/2$$

**Good**

$$\mathcal{P}_R \mathcal{P}^+ = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\mathcal{P}_L \mathcal{P}^+ = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\mathcal{P}_R \mathcal{P}^- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$\mathcal{P}_L \mathcal{P}^- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# The correlator as probability density matrix

---

$$\Phi\gamma^0 = \begin{pmatrix} f_1 & i\frac{(p_x+ip_y)}{M}h_1^\perp & 0 & 0 \\ -i\frac{(p_x-ip_y)}{M}h_1^\perp & f_1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

# The correlator as probability density matrix

---

$$\Phi\gamma^0 = \begin{pmatrix} f_1 & i\frac{(p_x+ip_y)}{M}h_1^\perp & 0 & 0 \\ -i\frac{(p_x-ip_y)}{M}h_1^\perp & f_1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

# The correlator as probability density matrix

$$\Phi\gamma^0 = \begin{pmatrix} R & | & L \\ | & f_1 & i\frac{(p_x+ip_y)}{M}h_1^\perp \\ L & | & R \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

The diagram shows a 3x3 matrix with boundary conditions. The top row has boundary conditions R (Right) and L (Left). The bottom row has boundary conditions L (Left) and R (Right). The diagonal elements are labeled  $f_1$ . The off-diagonal elements are labeled  $i\frac{(p_x+ip_y)}{M}h_1^\perp$ . The matrix is zero elsewhere.

# The correlator as probability density matrix

$$\Phi\gamma^0 = \begin{pmatrix} R & | & L \\ | & f_1 & i\frac{(p_x+ip_y)}{M}h_1^\perp \\ L & | & R \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

The diagram shows a 3x3 matrix with boundary conditions. The top row has boundary conditions R (Right) at both ends. The bottom row has boundary conditions L (Left) at both ends. The middle column has boundary conditions R (Right) at the top and L (Left) at the bottom. The matrix elements are: (1,1) =  $f_1$ , (1,2) =  $i\frac{(p_x+ip_y)}{M}h_1^\perp$ , (1,3) = 0, (2,1) =  $-i\frac{(p_x-ip_y)}{M}h_1^\perp$ , (2,2) =  $f_1$ , (2,3) = 0, (3,1) = 0, (3,2) = 0, (3,3) = 0.

- Probabilistic interpretation

# The correlator as probability density matrix

$$\Phi\gamma^0 = \begin{pmatrix} & R & L & & \\ R & f_1 & -i\frac{(p_x-ip_y)}{M}h_1^\perp & 0 & 0 \\ L & i\frac{(p_x+ip_y)}{M}h_1^\perp & f_1 & 0 & 0 \\ & 0 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 0 \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

Diagram illustrating the elements of the matrix:

- Top-left element:  $R$  (Right) vertex
- Top-right element:  $L$  (Left) vertex
- Bottom-left element:  $R$  (Right) vertex
- Bottom-right element:  $L$  (Left) vertex

- Probabilistic interpretation
- Positivity bounds

# The correlator as probability density matrix

$$\Phi\gamma^0 = \begin{pmatrix} R & | & R \\ | & f_1 & i\frac{(p_x+ip_y)}{M}h_1^\perp \\ L & | & L \\ | & -i\frac{(p_x-ip_y)}{M}h_1^\perp & f_1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

The diagram shows the correlation function  $\Phi\gamma^0$  represented as a 4x4 matrix. The rows and columns are labeled by two states: R and L. The matrix elements are:  
- Top-left element:  $f_1$   
- Top-right element:  $i\frac{(p_x+ip_y)}{M}h_1^\perp$   
- Bottom-left element:  $-i\frac{(p_x-ip_y)}{M}h_1^\perp$   
- Bottom-right element:  $f_1$   
- All other elements are zero.

- Probabilistic interpretation
- Positivity bounds
- Need of orbital angular momentum

# In $x$ -transversity basis

---

$$\Phi \gamma^0 = \begin{pmatrix} \text{Diagram: } \downarrow \text{ (yellow blob)} & \text{Diagram: } \uparrow \text{ (yellow blob)} \\ \text{Diagram: } \uparrow \text{ (yellow blob)} & \text{Diagram: } \downarrow \text{ (yellow blob)} \end{pmatrix} \sim \psi_i |P\rangle\langle P| \psi_j^\dagger$$

$$\begin{pmatrix} f_1 - \frac{p_y}{M} h_1^\perp & i \frac{p_x}{M} h_1^\perp \\ -i \frac{p_x}{M} h_1^\perp & f_1 + \frac{p_y}{M} h_1^\perp \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$$

# TMDs and their probabilistic interpretation

---

# TMDs and their probabilistic interpretation

---

quark pol.

	U	L	T
U	$f_1$		$h_1^\perp$
L		$g_{1L}$	$h_{1L}^\perp$
T	$f_{1T}^\perp$	$g_{1T}$	$h_1, h_{1T}^\perp$

Twist-2 TMDs

TMDs in black survive transverse-momentum integration  
TMDs in red are T-odd

# TMDs and their probabilistic interpretation

---


$$f_1 = \text{circle with blue center}$$


$$g_1 = \text{circle with black dot} - \text{circle with red cross}$$

$$h_1 = \text{two circles connected by a horizontal line with arrows pointing right}$$


# TMDs and their probabilistic interpretation


---


$$f_{1T}^\perp = - \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


$$f_1 = \begin{array}{c} \text{---} \\ \text{---} \end{array}$$



$$h_1^\perp = - \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


$$g_1 = \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


$$g_{1T} = - \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


$$h_1 = \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


$$h_{1L}^\perp = - \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


$$h_{1T}^\perp = - \begin{array}{c} \text{---} \\ \text{---} \end{array}$$


# A short note on $g_T$

---

$$\begin{aligned}\Phi(x, S_x) \gamma^0 &= \frac{1}{2} \left\{ f_1 \gamma^- + S_L g_1 \gamma_5 \gamma^- - h_1 S_T \gamma_5 \gamma^1 \gamma^- \right\} \gamma^0 \\ &\quad + \frac{M}{2P^+} \left\{ e - g_T S_T \gamma_5 \gamma^1 + h_L S_L \frac{[\gamma^-, \gamma^+] \gamma_5}{2} \right\} \gamma^0\end{aligned}$$

# A short note on $g_T$

---

$$\begin{aligned}\Phi(x, S_x) \gamma^0 &= \frac{1}{2} \left\{ f_1 \gamma^- + S_L g_1 \gamma_5 \gamma^- - h_1 S_T \gamma_5 \gamma^1 \gamma^- \right\} \gamma^0 \\ &\quad + \frac{M}{2P^+} \left\{ e - g_T S_T \gamma_5 \gamma^1 + h_L S_L \frac{[\gamma^-, \gamma^+] \gamma_5}{2} \right\} \gamma^0\end{aligned}$$

$$\gamma_5 \gamma^- \gamma^0 = \sqrt{2} \begin{pmatrix} \sigma^3 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^1 \gamma^- \gamma^0 = \sqrt{2} \begin{pmatrix} \sigma^1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^1 \gamma^0 = \sqrt{2} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}$$

# A short note on $g_T$

---

$$\begin{aligned}\Phi(x, S_x) \gamma^0 &= \frac{1}{2} \left\{ f_1 \gamma^- + S_L g_1 \gamma_5 \gamma^- - h_1 S_T \gamma_5 \gamma^1 \gamma^- \right\} \gamma^0 \\ &\quad + \frac{M}{2P^+} \left\{ e - g_T S_T \gamma_5 \gamma^1 + h_L S_L \frac{[\gamma^-, \gamma^+] \gamma_5}{2} \right\} \gamma^0\end{aligned}$$

$$\gamma_5 \gamma^- \gamma^0 = \sqrt{2} \begin{pmatrix} \sigma^3 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^1 \gamma^- \gamma^0 = \sqrt{2} \begin{pmatrix} \sigma^1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^1 \gamma^0 = \sqrt{2} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}$$

My impression is that it's relevant for the Bakker-Trueman-Leader transverse angular momentum sum-rule

# A short note on $g_T$

---

$$\begin{aligned}\Phi(x, S_x) \gamma^0 &= \frac{1}{2} \left\{ f_1 \gamma^- + S_L g_1 \gamma_5 \gamma^- - h_1 S_T \gamma_5 \gamma^1 \gamma^- \right\} \gamma^0 \\ &\quad + \frac{M}{2P^+} \left\{ e - g_T S_T \gamma_5 \gamma^1 + h_L S_L \frac{[\gamma^-, \gamma^+] \gamma_5}{2} \right\} \gamma^0\end{aligned}$$

$$\gamma_5 \gamma^- \gamma^0 = \sqrt{2} \begin{pmatrix} \sigma^3 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^1 \gamma^- \gamma^0 = \sqrt{2} \begin{pmatrix} \sigma^1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^1 \gamma^0 = \sqrt{2} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix}$$

My impression is that it's relevant for the Bakker-Trueman-Leader transverse angular momentum sum-rule

<http://www.ts.infn.it/eventi/transversitySR/>

# Unpolarized sector

---

$$F_{UU,T} = \mathcal{C}[f_1 D_1],$$

$$F_{UU,L} = \mathcal{O}\left(\frac{M^2}{Q^2}, \frac{q_T^2}{Q^2}\right),$$

$$F_{UU}^{\cos\phi_h} = \frac{2M}{Q} \mathcal{C} \left[ -\frac{\hat{\boldsymbol{h}} \cdot \boldsymbol{k}_T}{M_h} \left( x h H_1^\perp + \frac{M_h}{M} f_1 \frac{\tilde{D}^\perp}{z} \right) - \frac{\hat{\boldsymbol{h}} \cdot \boldsymbol{p}_T}{M} \left( x f^\perp D_1 + \frac{M_h}{M} h_1^\perp \frac{\tilde{H}}{z} \right) \right],$$

$$F_{UU}^{\cos 2\phi_h} = \mathcal{C} \left[ -\frac{2 \left( \hat{\boldsymbol{h}} \cdot \boldsymbol{k}_T \right) \left( \hat{\boldsymbol{h}} \cdot \boldsymbol{p}_T \right) - \boldsymbol{k}_T \cdot \boldsymbol{p}_T}{MM_h} h_1^\perp H_1^\perp \right],$$

$$\mathcal{C}[wfD] = \sum_a x e_a^2 \int d^2 \boldsymbol{p}_T d^2 \boldsymbol{k}_T \delta^{(2)}(\boldsymbol{p}_T - \boldsymbol{k}_T - \boldsymbol{P}_{h\perp}/z) w(\boldsymbol{p}_T, \boldsymbol{k}_T) f^a(x, p_T^2) D^a(z, k_T^2),$$

# Longitudinally polarized beam or/and target

---

$$F_{LU}^{\sin \phi_h} = \frac{2M}{Q} \mathcal{C} \left[ -\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M_h} \left( xe H_1^\perp + \frac{M_h}{M} f_1 \frac{\tilde{G}^\perp}{z} \right) + \frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} \left( x g^\perp D_1 + \frac{M_h}{M} h_1^\perp \frac{\tilde{E}}{z} \right) \right],$$

$$F_{UL}^{\sin \phi_h} = \frac{2M}{Q} \mathcal{C} \left[ -\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M_h} \left( x h_L H_1^\perp + \frac{M_h}{M} g_{1L} \frac{\tilde{G}^\perp}{z} \right) + \frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} \left( x f_L^\perp D_1 - \frac{M_h}{M} h_{1L}^\perp \frac{\tilde{H}}{z} \right) \right],$$

$$F_{UL}^{\sin 2\phi_h} = \mathcal{C} \left[ -\frac{2 (\hat{\mathbf{h}} \cdot \mathbf{k}_T) (\hat{\mathbf{h}} \cdot \mathbf{p}_T) - \mathbf{k}_T \cdot \mathbf{p}_T}{MM_h} h_{1L}^\perp H_1^\perp \right],$$

$$F_{LL} = \mathcal{C} [g_{1L} D_1],$$

$$F_{LL}^{\cos \phi_h} = \frac{2M}{Q} \mathcal{C} \left[ \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M_h} \left( xe_L H_1^\perp - \frac{M_h}{M} g_{1L} \frac{\tilde{D}^\perp}{z} \right) - \frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} \left( x g_L^\perp D_1 + \frac{M_h}{M} h_{1L}^\perp \frac{\tilde{E}}{z} \right) \right],$$

# Transversely polarized beam

---

$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = \mathcal{C} \left[ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} f_{1T}^\perp D_1 \right],$$

$$F_{UT,L}^{\sin(\phi_h - \phi_S)} = \mathcal{O} \left( \frac{M^2}{Q^2}, \frac{q_T^2}{Q^2} \right),$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = \mathcal{C} \left[ -\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M_h} h_1 H_1^\perp \right],$$

$$F_{UT}^{\sin(3\phi_h - \phi_S)} = \mathcal{C} \left[ \frac{2 (\hat{\mathbf{h}} \cdot \mathbf{p}_T) (\mathbf{p}_T \cdot \mathbf{k}_T) + \mathbf{p}_T^2 (\hat{\mathbf{h}} \cdot \mathbf{k}_T) - 4 (\hat{\mathbf{h}} \cdot \mathbf{p}_T)^2 (\hat{\mathbf{h}} \cdot \mathbf{k}_T)}{2M^2 M_h} h_{1T}^\perp H_1^\perp \right],$$

$$\begin{aligned} F_{UT}^{\sin \phi_S} &= \frac{2M}{Q} \mathcal{C} \left\{ \left( x f_T D_1 - \frac{M_h}{M} h_1 \frac{\tilde{H}}{z} \right) \right. \\ &\quad \left. - \frac{\mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[ \left( x h_T H_1^\perp + \frac{M_h}{M} g_{1T} \frac{\tilde{G}^\perp}{z} \right) - \left( x h_T^\perp H_1^\perp - \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{D}^\perp}{z} \right) \right] \right\}, \end{aligned}$$

$$\begin{aligned} F_{UT}^{\sin(2\phi_h - \phi_S)} &= \frac{2M}{Q} \mathcal{C} \left\{ \frac{2 (\hat{\mathbf{h}} \cdot \mathbf{p}_T)^2 - \mathbf{p}_T^2}{2M^2} \left( x f_T^\perp D_1 - \frac{M_h}{M} h_{1T}^\perp \frac{\tilde{H}}{z} \right) \right. \\ &\quad \left. - \frac{2 (\hat{\mathbf{h}} \cdot \mathbf{k}_T) (\hat{\mathbf{h}} \cdot \mathbf{p}_T) - \mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[ \left( x h_T H_1^\perp + \frac{M_h}{M} g_{1T} \frac{\tilde{G}^\perp}{z} \right) + \left( x h_T^\perp H_1^\perp - \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{D}^\perp}{z} \right) \right] \right\}, \end{aligned}$$

# Trasversely pol. target and long. pol. beam

---

$$F_{LT}^{\cos(\phi_h - \phi_S)} = \mathcal{C} \left[ \frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} g_{1T} D_1 \right],$$

$$\begin{aligned} F_{LT}^{\cos \phi_S} &= \frac{2M}{Q} \mathcal{C} \left\{ - \left( xg_T D_1 + \frac{M_h}{M} h_1 \frac{\tilde{E}}{z} \right) \right. \\ &\quad \left. + \frac{\mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[ \left( xe_T H_1^\perp - \frac{M_h}{M} g_{1T} \frac{\tilde{D}^\perp}{z} \right) + \left( xe_T^\perp H_1^\perp + \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{G}^\perp}{z} \right) \right] \right\}, \end{aligned}$$

$$\begin{aligned} F_{LT}^{\cos(2\phi_h - \phi_S)} &= \frac{2M}{Q} \mathcal{C} \left\{ - \frac{2(\hat{\mathbf{h}} \cdot \mathbf{p}_T)^2 - \mathbf{p}_T^2}{2M^2} \left( xg_T^\perp D_1 + \frac{M_h}{M} h_{1T}^\perp \frac{\tilde{E}}{z} \right) \right. \\ &\quad + \frac{2(\hat{\mathbf{h}} \cdot \mathbf{k}_T)(\hat{\mathbf{h}} \cdot \mathbf{p}_T) - \mathbf{k}_T \cdot \mathbf{p}_T}{2MM_h} \left[ \left( xe_T H_1^\perp - \frac{M_h}{M} g_{1T} \frac{\tilde{D}^\perp}{z} \right) \right. \\ &\quad \left. \left. - \left( xe_T^\perp H_1^\perp + \frac{M_h}{M} f_{1T}^\perp \frac{\tilde{G}^\perp}{z} \right) \right] \right\} \end{aligned}$$

# The gauge link

---

# Need of a gauge link

---

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4\xi \ e^{ip\cdot\xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle$$

# Need of a gauge link

---

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4\xi \ e^{ip\cdot\xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle$$

$$\psi(\xi) \rightarrow e^{i\alpha(\xi)} \psi(\xi)$$

# Need of a gauge link

---

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4\xi \ e^{ip\cdot\xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle$$

$$\psi(\xi) \rightarrow e^{i\alpha(\xi)} \psi(\xi)$$

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4\xi \ e^{ip\cdot\xi} \langle P, S | \bar{\psi}_j(0) U_{[0,\xi]} \psi_i(\xi) | P, S \rangle$$

# Need of a gauge link

---

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4\xi \ e^{ip\cdot\xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle$$

$$\psi(\xi) \rightarrow e^{i\alpha(\xi)} \psi(\xi)$$

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4\xi \ e^{ip\cdot\xi} \langle P, S | \bar{\psi}_j(0) U_{[0,\xi]} \psi_i(\xi) | P, S \rangle$$

$$U(\xi_1, \xi_2) \rightarrow e^{i\alpha(\xi_1)} U(\xi_1, \xi_2) e^{-i\alpha(\xi_2)}.$$

# Need of a gauge link

---

$$\Phi_{ij}(p,P,S) = \frac{1}{(2\pi)^4} \int d^4\xi \; e^{ip\cdot\xi} \langle P,S | \bar{\psi}_j(0) \psi_i(\xi) | P,S \rangle$$

$$\psi(\xi)\rightarrow e^{i\alpha(\xi)}\,\psi(\xi)$$


$$\Phi_{ij}(p,P,S) = \frac{1}{(2\pi)^4} \int d^4\xi \; e^{ip\cdot\xi} \langle P,S | \bar{\psi}_j(0) U_{[0,\xi]} \psi_i(\xi) | P,S \rangle$$

$$U(\xi_1,\xi_2)\rightarrow e^{i\alpha(\xi_1)}\,U(\xi_1,\xi_2)\,e^{-i\alpha(\xi_2)}.$$

$$U_{[a,b]}=\mathcal{P}\exp\left[-ig\int_a^b d\eta^\mu A_\mu(\eta)\right]$$

# Birth of the gauge link

---




(a)

$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\alpha \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \gamma_\nu g A^\alpha(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0}$$

# Birth of the gauge link

---




(a)

$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\alpha \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \gamma_\nu g A^\alpha(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0}$$

$$i \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \approx i \frac{k^- \gamma^+}{-2l^+ k^- + i\epsilon} \approx \frac{i}{2} \frac{\gamma^+}{-l^+ + i\epsilon}$$

# Birth of the gauge link

---



(a)


$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\alpha \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \gamma_\nu g A^\alpha(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0}$$

$$i \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \approx i \frac{k^- \gamma^+}{-2l^+ k^- + i\epsilon} \approx \frac{i}{2} \frac{\gamma^+}{-l^+ + i\epsilon}$$

$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu + \frac{\gamma_- + \gamma_+}{2} \gamma_\nu (ig) \frac{A^+(\ )}{-l^+ + i} (\ ) | P, S \rangle \Big|_{\eta^+ = 0}$$

# Birth of the gauge link

---



(a)

$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\alpha \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \gamma_\nu g A^\alpha(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0}$$


$$i \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \approx i \frac{k^- \gamma^+}{-2l^+ k^- + i\epsilon} \approx \frac{i}{2} \frac{\gamma^+}{-l^+ + i\epsilon}$$

$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\nu (ig) \frac{A^+(\eta)}{-l^+ + i} (\xi) | P, S \rangle \Big|_{\eta^+ = 0}$$


$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\nu (-ig) \int_{\infty^-}^{\xi^-} d\eta^- A^+(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0; \eta_T = \xi_T}$$

# Birth of the gauge link

---



$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\nu (-ig) \int_{\infty^-}^{\xi^-} d\eta^- A^+(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+=0; \eta_T = \xi_T}$$




$$2MW^{\mu\nu}(q, P, S) \approx \sum_q e_q^2 \frac{1}{2} \text{Tr} [\Phi(x_B, S) \gamma^\mu \gamma^+ \gamma^\nu].$$


$$\Phi^{(a)}(x, S) \sim \langle P, S | \bar{\psi}(0) (-ig) \int_{\infty^-}^{\xi^-} d\eta^- A^+(\eta) \psi(\xi) | P, S \rangle$$

# Birth of the gauge link

---




$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\nu (-ig) \int_{\infty^-}^{\xi^-} d\eta^- A^+(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+=0; \eta_T = \xi_T}$$



$$2MW^{\mu\nu}(q, P, S) \approx \sum_q e_q^2 \frac{1}{2} \text{Tr} [\Phi(x_B, S) \gamma^\mu \gamma^+ \gamma^\nu].$$



$$\Phi^{(a)}(x, S) \sim \langle P, S | \bar{\psi}(0) (-ig) \int_{\infty^-}^{\xi^-} d\eta^- A^+(\eta) \psi(\xi) | P, S \rangle$$



# Shape of the gauge link

---




$$\Phi(x, S) \sim \langle P, S | \bar{\psi}(0) U_{[0, \infty^-]} U_{[\infty^-, \xi^-]} \psi(\xi) | P, S \rangle$$



# Shape of the gauge link


---

$$\Phi(x, S) \sim \langle P, S | \bar{\psi}(0) U_{[0, \infty^-]} U_{[\infty^-, \xi^-]} \psi(\xi) | P, S \rangle$$



# Gauge link in Drell-Yan

---



$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\alpha \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \gamma_\nu g A^\alpha(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0}$$

$$i \frac{k - l + m}{(k - l)^2 - m^2 + i\epsilon} \approx i \frac{-(-k)^- \gamma^+}{2l^+(-k)^- + i\epsilon} \approx \frac{i}{2} \frac{\gamma^+}{-l^+ - i\epsilon}$$

$$2MW_{\mu\nu}^{(a)} \sim \langle P, S | \bar{\psi}(0) \gamma_\mu \gamma^+ \gamma_\nu (-ig) \int_{-\infty^-}^{\xi^-} d\eta^- A^+(\eta) \psi(\xi) | P, S \rangle \Big|_{\eta^+ = 0; \eta_T = \xi_T}$$

# Gauge link for TMDs

---


$$\Phi_{ij}(x, \textcolor{red}{p_T}) = \int \frac{d\xi^- d^2\xi_T}{8\pi^3} e^{ip \cdot \xi} \langle P | \bar{\psi}_j(0) \textcolor{red}{U}_{[0,\xi]} \psi_i(\xi) | P \rangle \Big|_{\xi^+ = 0}$$

# Gauge link for TMDs

---

$$\Phi_{ij}(x, \textcolor{red}{p_T}) = \int \frac{d\xi^- d^2\xi_T}{8\pi^3} e^{ip \cdot \xi} \langle P | \bar{\psi}_j(0) \textcolor{red}{U}_{[0,\xi]} \psi_i(\xi) | P \rangle \Big|_{\xi^+ = 0}$$

SIDIS




# Gauge link for TMDs


---

$$\Phi_{ij}(x, \textcolor{red}{p_T}) = \int \frac{d\xi^- d^2\xi_T}{8\pi^3} e^{ip \cdot \xi} \langle P | \bar{\psi}_j(0) \textcolor{red}{U}_{[0,\xi]} \psi_i(\xi) | P \rangle \Big|_{\xi^+ = 0}$$

SIDIS




Drell-Yan




# Gauge link for TMDs

$$\Phi_{ij}(x, \textcolor{red}{p_T}) = \int \frac{d\xi^- d^2\xi_T}{8\pi^3} e^{ip \cdot \xi} \langle P | \bar{\psi}_j(0) \textcolor{red}{U}_{[0,\xi]} \psi_i(\xi) | P \rangle \Big|_{\xi^+ = 0}$$

SIDIS



Drell-Yan



$pp$  to hadrons

